

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
13 June 2002 (13.06.2002)

PCT

(10) International Publication Number  
**WO 02/46746 A2**

(51) International Patent Classification: **G01N 33/48**

Tyne & Wear NE35 9PD (GB). **GARDNER, Michael, James** [GB/GB]; Immunodiagnostic Systems Ltd, Boldon Business Park, Boldon, Tyne & Wear NE35 9PD (GB).

(21) International Application Number: **PCT/GB01/05395**

(22) International Filing Date: 6 December 2001 (06.12.2001)

(74) Agents: **DAVIES, Punita et al.**; Kilburn & Strode, 20 Red Lion Street, London WC1R 4PJ (GB).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0029729.1 6 December 2000 (06.12.2000) GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): **IMMUNODIAGNOSTIC SYSTEMS LTD** [GB/GB]; Boldon Business Park, Boldon, Tyne & Wear NE35 9PD (GB).


(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **LAURIE, David** [GB/GB]; Immunodiagnostic Systems Ltd, Boldon Business Park, Boldon, Tyne & Wear NE35 9PD (GB). **BARNES, Alexander, Kirkley** [GB/GB]; Immunodiagnostic Systems Ltd, Boldon Business Park, Boldon,

[Continued on next page]

(54) Title: METHOD FOR DETECTION OF VITAMIN D METABOLITES



(57) Abstract: The present invention relates to a method for measuring the amount of vitamin D metabolite in a sample of plasma or serum, comprising the step of (a) adding to a serum or plasma sample a non-competitive displacement agent to effect separation of any vitamin D metabolite in the sample from protein to which it is bound, such that any vitamin D metabolite may be detected and/or measured; and (b) detecting or measuring the amount of vitamin D metabolite in the sample of (a). Such methods are useful in determining the vitamin D status of the subject, for example in the diagnosis of disease. Also provided are agents and kits for use in performing the methods of the invention.



WO 02/46746 A2



**Published:**

— without international search report and to be republished  
upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

## METHOD FOR DETECTION OF VITAMIN D METABOLITES

The present invention relates to a method for measuring the amount of a vitamin D metabolite present in a sample of plasma or serum, and in particular to a method which enables measurement of the amount of a vitamin D metabolite in a sample of plasma or serum without the need for prior extraction from the sample of vitamin D metabolite binding proteins. The present invention also relates to methods for determining the vitamin D status of a subject, for use in the diagnosis of disease, and to agents and kits for use in performing the methods of the invention.

10

Vitamin D status is an indication of the level of circulating vitamin D or vitamin D metabolites in a subject, and is an important factor in the cause of a number of diseases including rickets in children and osteomalacia in adults. It is also useful in identifying the underlying causes of hypercalcaemia and hypocalcaemia. Vitamin D and its metabolism are the subject of great clinical interest, and in line with this there has been a steady increase in the effort towards improving methods for the measurement of vitamin D and its metabolites in body fluids.

Vitamin D is available to man as either vitamin D<sub>2</sub> or vitamin D<sub>3</sub>. Vitamin D<sub>2</sub> is produced outside the body by irradiation of ergosterol from yeast and fungi, and is found in man when taken in the form of fortified foods or pharmaceutical preparations. Vitamin D<sub>3</sub>, on the other hand, is formed in animals from 7-dehydrocholesterol upon exposure to *uv* light. In man, this reaction occurs in the skin. Vitamin D<sub>3</sub> is also available in the diet, for example from fish liver oils. Vitamin D, in the form of vitamin D<sub>2</sub> or D<sub>3</sub>, is rapidly converted to the circulating metabolite, 25-hydroxy vitamin D, which is found outside cells, tightly bound to circulating vitamin D binding protein (DBP). Due to the rapid conversion of vitamin D to its' metabolite, measurement of vitamin D does not give a useful indication of the vitamin D status of a subject. Other metabolites of vitamin D, such

20

25

as  $1\alpha,25$ -dihydroxy vitamin D, circulate at a concentration 1000 times lower than non- $1\alpha$ -hydroxylated metabolites such as 25-hydroxy vitamin D, and so do not contribute significantly to the estimation of total circulating vitamin D metabolite. For this reason, the  $1\alpha$ -hydroxylated metabolites do not provide a direct or useful indication of vitamin D status. 25-hydroxy vitamin D is the metabolite with the highest serum concentration, and is easy to measure. It has therefore become the most common marker of vitamin D status in a subject.

The established methods for measurement of 25-hydroxy vitamin D provide a result which includes contributions from other vitamin D hydroxylated metabolites, principally 24,25-dihydroxy vitamin D but also 25,26-dihydroxy vitamin D and other metabolites. These other hydroxylated metabolites contribute 1 to 7% of the measure of total circulating vitamin D metabolites, the remainder being 25-hydroxy vitamin D. Whilst for many applications, including research, it is sufficient to measure the total amount of circulating vitamin D metabolites, for other applications separation of these metabolites is required to allow specific measurement of 25-hydroxy vitamin D. It is generally accepted that measurement of 25-hydroxy vitamin D levels in serum or plasma samples is the preferred method of determining the vitamin D status of a subject.

25-hydroxy vitamin D is found circulating in the serum or plasma, tightly bound to DBP (also known as Gc globulin). It is hydrophobic, and in addition to binding to DBP, it is found to bind strongly to other binding proteins such as albumin. The presence of such binding proteins, in particular DBP, presents a major difficulty in measuring serum or plasma levels of 25-hydroxy vitamin D.

To deal with this problem, methods for measurement of 25-hydroxy vitamin D to date have relied upon removal of DBP or other binding proteins from the sample to be analysed. This has typically involved an extraction step which separates the DBP

from the vitamin D metabolite and removes the DBP from the sample, leaving behind the vitamin D metabolite. Extraction has been achieved by a number of methods, including solvent based extraction by adding to the sample an organic solvent such as chloroform, hexane or ethyl acetate and hexane. The organic and aqueous layers are separated and the solvent evaporated. The residue is then reconstituted in a water miscible solvent such as ethanol. Reverse phase cartridge extraction methods have also been used. Other traditional methods include the use of HPLC and chromatography to achieve separation of individual vitamin D metabolites and exclude from the sample interfering factors such as binding proteins.

10

Simplification of the methods for extraction and separation has been a key feature in the improvement of assays for vitamin D and its metabolites. For example, the chromatography step has been eliminated by the use of water miscible solvent such as acetonitrile to precipitate binding proteins. The binding proteins are then separated from the supernatant by centrifugation, and the supernatant is then analysed by immunoassay or competitive protein binding assay to determine the amount of vitamin D metabolite present.

15

While removal of vitamin D binding proteins from serum samples by organic solvents is effective, the process requires evaporation of volatile organic solvents which is not attractive for routine use in clinical biochemical laboratories. Although the use of water miscible solvents such as acetonitrile has largely eliminated the need for solvent evaporation, these have the disadvantage of being toxic. In addition, the requirement for a separate extraction step inevitably lengthens the assay time.

20

25

Boullion *et al* (Clin. Chem. 30/11 1731-1736 (1984)) describe "Two Direct (Non-Chromatographic) Assays for 25-Hydroxy Vitamin D". Whilst the assays described do not require a chromatography step as required in more traditional methods, they still require extraction of the vitamin D binding proteins from the sample by the use of

solvent precipitation. Thus, the assays described in this publication suffer from the drawbacks mentioned above.

Holick *et al* (5,981,779) describe methods for assaying vitamin D and its' metabolites in milk and other biological fluids. One of the solid phase assays described uses a competitive binding assay to measure the vitamin D metabolite in the sample, and comprises the steps of: (a) providing a solid phase support having immobilised thereon a protein or antibody which is capable of binding labelled vitamin D; (b) contacting the solid phase with a solution of labelled vitamin D to allow it to bind to the solid phase; (c) washing the solid phase to remove unbound labelled vitamin D; (d) contacting the solid phase with a sample suspected of containing vitamin D or its metabolite, for a time sufficient to displace the labelled compound; (e) collecting the liquid of (d); and (f) measuring the amount of labelled vitamin D in the liquid, which is proportional to the amount of vitamin D or metabolite in the test sample. This method is shown as a schematic in Figure 3, when performed using milk as the test sample.

The method described is essentially a "free hormone" assay. In the example illustrated, the sample is milk which contains little DBP. The hydrophobic vitamin D will be bound to other milk proteins, the binding of which will be relatively low affinity compared with the binding for DBP. This is the basis on which a suitable competition between sample-derived vitamin D and labelled vitamin D tracer is achieved in the assay for milk samples. In samples containing significant amounts of DBP, for example serum, where the DBP level is ~400mg/L, approximately 6% of the  $\alpha$ -globulin fraction of serum, the "free" (i.e. that not bound to DBP) vitamin D is a very small proportion of the total (0.04% for serum). Such an assay is not suitable for measurement of vitamin D metabolite in these samples, a view which is further supported by Holick *et al* in a recent publication describing by a variation of the illustrated assay for application to serum samples. However, the only significant

difference between the illustrated milk assay and the serum assay is the need for an additional solvent extraction stage to remove DBP and other proteins from the sample (11<sup>th</sup> Workshop on Vitamin D, 27 May-1 June 2000, Nashville, USA). If serum was substituted for milk in the original assay, the labelled vitamin D originally bound to the DBP on the tube would simply equilibrate with the DBP present in the sample, of which 98% have unoccupied D binding sites. Such an assay would therefore only provide a measure of the capacity of the sample to bind a vitamin D-analogue, and not the sample's vitamin D content.

10 The Holick patent teaches that measurement of vitamin D (or metabolites) is possible in milk samples without need for an extraction process. It does not teach efficient methods by which the difficulties of vitamin D measurement in serum can be overcome.

15 Measurement of steroid hormones in serum, plasma or other biological fluids is carried out by a direct immunoassay, without the need for a separate extraction procedure. This has been achieved by the use of a steroid analogue which binds to the relevant steroid binding proteins, but does not cross react with the antibody used in the immunoassay. The steroid analogue saturates the steroid binding protein, displacing  
20 the steroid and allowing the steroid to bind to the antibodies of the immunoassay.

Use of a (specific) competitive displacer such as a vitamin D analogue that does not cross-react with the assay antibody, should be able to provide a "direct assay" (by analogy to direct steroid measurement methods). However there are two difficulties.  
25 Firstly, the concentration of DBP is very high in serum samples, typically ~400mg/L (~7 $\mu$ mol/l), and typically only ~2% of DBP has 25 hydroxy-vitamin D bound, the remaining ~98% DBP remains unoccupied and will bind potential competitive displacers. As a result the concentration of a specific competitive displacer must be sufficient to saturate the DBP and would need to be in excess if the affinity was lower

than that of 25 hydroxyvitamin D. Vitamin D analogues are particularly expensive due to the complexity of vitamin D chemistry. Secondly, the specificity of DBP (for natural vitamin D metabolites) means that even with the availability of large numbers of different vitamin D analogues, very few have affinity for DBP that can compete with 25 hydroxyvitamin D.

The present invention aims to overcome or ameliorate the problems associated with the prior art, by providing an improved method for measuring vitamin D metabolite present in a sample of serum or plasma.

10

Thus, in a first aspect of the present invention there is provided a method for measuring vitamin D metabolite present in a plasma or serum sample of a subject, the method comprising:

- (a) adding to a sample of serum or plasma a non-competitive displacement agent to effect separation of any vitamin D metabolite in the sample from protein to which it is bound, such that any vitamin D metabolite can be detected and/or measured; and
- (b) detecting or measuring the amount of vitamin D metabolite in the sample of (a).

20

Thus, the present invention satisfies the pressing need for a simple yet effective method for measuring vitamin D metabolite in a serum or plasma sample. It is based upon the surprising discovery of non-competitive displacement agents which enable effective separation of vitamin D metabolites from binding proteins to enable the amount of vitamin D metabolites to be detected or measured, without competing with the protein or requiring its extraction from the sample. In essence, the invention provides for the first time displacement agents for use in vitamin D metabolite analysis which are not analogues or competitors of the vitamin D metabolites to be measured. The elimination of the need of both the extraction step and competitive displacement

25



agents has enabled a method which is more efficient and cost effective than previous methods, and thus more suited to routine use in clinical biochemical laboratories or personal use.

5 The present invention may be performed on any sample of plasma or serum, preferably from a subject. The subject whose plasma or serum is to be analyzed may be one for whom it is desirable to determine vitamin D status. Preferably, the subject may be a mammal. More preferably, the subject is a human, and most preferably a child.

10

Measuring vitamin D metabolite present in a sample of plasma or serum may include both detecting the presence of any vitamin D metabolite in the sample, or more preferably, determining the amount of vitamin D metabolite present. The amount may be compared with a key detailing whether the amount represents a deficiency or  
15 excess of vitamin D metabolite.

Any one or more metabolites of vitamin D may be measured in the method of the present invention. In a preferred embodiment, a specific vitamin D metabolite is measured in a sample, although it is envisaged that for some applications it may be  
20 preferred to measure two or more types of metabolite present. Examples of metabolites include 25-hydroxy vitamin D<sub>2</sub>, 25-hydroxy vitamin D<sub>3</sub>, 24,25-dihydroxy vitamin D<sub>3</sub> and 25,26-dihydroxy vitamin D. 25-hydroxy vitamins D<sub>2</sub> or D<sub>3</sub>, or analogues thereof, are the preferred metabolites to be measured in the method of the invention.

25

The method of the invention is not restricted to the separation, or displacement, of a vitamin D metabolite from DBP, but may be used in the separation of a vitamin D metabolite from any factor to which it is bound. Typically, such factors will be proteins, for example, serum albumin.

30

By displacement is meant full or partial separation of some or all of the vitamin D metabolite from the factor to which it is bound in the sample. Preferably, the displacement of the vitamin D metabolite from the binding factor is sufficient to enable the vitamin D metabolite to be detected and/or measured, for example in a binding assay. It is preferred that substantially all of the vitamin D metabolite present in the sample is sufficiently displaced from the binding factor, to enable it to be measured. In this context, "substantially" means at least 95%, 98% and preferably at least 99% of the vitamin D metabolite is displaced. This ensures accuracy of a test based upon the method of the invention.

Any non-competitive agent capable of achieving displacement, or separation, of the vitamin D metabolite from the binding protein may be used in step (a) of the method of the invention. Preferred agents for use in the present invention are chemical reagents which may act by disrupting or destroying the bond between vitamin D metabolite and binding factor. Preferred chemical reagents for use as displacement agents are those which bind to vitamin D binding proteins such as albumin with sufficient affinity to assist displacement of the vitamin D metabolite. In a most preferred embodiment of the present invention, the displacement agent is a chemical reagent comprising, preferably in a buffer, 8-anilino-1-naphthalenesulphonic acid ammonium salt, 3-(acetylbenzyl)-4-hydroxycoumarin and a water miscible solvent. Any suitable buffer may be used, for example phosphate buffered saline. Suitable water miscible solvents will be known to persons skilled in the art and include ethanol, acetonitrile, propan-1-ol, propan-2-ol, acetone, dimethylformamide and methyl sulphoxide. The preferred water miscible solvent is methanol. The suitable concentrations and proportions of the three reagents for producing the displacement agent can be readily ascertained by persons skilled in the art. Preferred concentrations of the reagents are 0.5 to 10g/l, more preferably 1.60g/l 8-anilino-1-naphthalenesulphonoc acid ammonium salt; 50 to 1000mg/l, and more preferably 160mg/l 3-(acetylbenzyl)-4-hydroxycoumarin and 10 to 300ml/l, more preferably 160ml/l water miscible solvent such as methanol.

Once the vitamin D metabolite has been displaced from the binding factor in the sample, the presence or amount of vitamin D metabolite in the sample may be determined. Any suitable method may be used for this purpose, and such methods will  
5 be familiar to persons skilled in the art. Examples of suitable methods include binding assays, such as competitive or non-competitive binding assays, immunoassays, or direct labeling of unbound vitamin D metabolites.

The step (b) of measuring the amount of vitamin D metabolite in the sample may be  
10 performed subsequent to, or simultaneously with step (a). Preferably, the step (b) is carried out simultaneously with step (a), thus providing a single step assay or measurement of vitamin D metabolite in a serum or plasma sample.

One of the preferred methods for measuring displaced vitamin D metabolite in a  
15 sample is by way of a competitive binding assay. Suitable competitive binding assays take various forms, and will be well known to persons skilled in the art. A typical competitive binding assay will comprise contacting a receptor with a labeled form of a ligand and a sample suspected of containing an unlabelled form of the same ligand. The amount of labeled ligand which is found bound to the receptor is indicative of the  
20 proportion of unlabeled ligand in the sample. Alternatively, the competitive binding assay may comprise providing receptors bound to a labeled form of the ligand, adding to the receptors the sample suspected of containing the unlabelled form of the ligand, and measuring the amount of displaced labeled ligand which is indicative of the amount of unlabelled ligand present.

25

In a preferred embodiment of the first aspect of the invention, there is provided a method of measuring the amount of displaced vitamin D metabolite in a sample, comprising the steps of:

(a) providing a support having immobilised thereon a binding factor capable of  
30 binding a vitamin D metabolite;

(b) contacting the support with a sample comprising the vitamin D metabolite to be measured;

(c) contacting the support with a labeled form of the vitamin D metabolite;

(d) measuring the amount of labeled vitamin D metabolite left bound to the support,

wherein the amount of labeled vitamin D metabolite bound to the support is proportional to the amount of vitamin D metabolite in the sample.

Where it is preferred to perform the binding assay simultaneously with the displacement step, the method of the first aspect may comprise the steps of

(a) providing a support having immobilised thereon a binding factor capable of binding a vitamin D metabolite;

(b) contacting the support with a serum or plasma sample of a subject;

(c) adding to the support a non-competitive displacement agent to effect separation of the vitamin D metabolite from protein to which it is bound;

(d) contacting the support with a labelled form of the vitamin D metabolite;

(e) measuring the amount of labelled vitamin D metabolite left bound to the support;

wherein the amount of labelled vitamin D metabolite bound to the support is proportional to the amount of vitamin D metabolite in the sample.

The binding factor to be immobilized on the support may be any factor capable of binding a labelled or unlabelled form of the vitamin D metabolite. The binding factor may be a protein, such as albumin or DBP, or an antibody specific for the particular vitamin D metabolite of interest. In a most preferred embodiment, the binding factor is an antibody against the vitamin D metabolite. Preferably, the antibody is polyclonal.

In an alternative embodiment, the vitamin D metabolite may be measured by way of an immunoassay. Any suitable form of immunoassay may be used, and these will be

known to persons skilled in the art. For example, the displaced vitamin D may be captured on a support of vitamin D metabolite specific antibodies, and then quantified using a second, labeled vitamin D metabolite specific antibody.

- 5 Any suitable labeling means may be used in the above methods. Suitable labels include enzymatic labels, such as alkaline phosphatase, peroxidase, biotin-streptavidin, fluorescent labels such as fluorescein or rhodamine, or chemiluminescent labels such as luminol, acridium esters, and 1,2-dioxetanes. Also included are combinations of the above, for example avidin-fluorescein, as an alternative for avidin-HRP.

10

In a second aspect of the invention, there is provided a method for determining the vitamin D status of a subject, the method comprising:

- (a) adding to a sample of serum or plasma a non-competitive displacement agent to effect separation of any vitamin D metabolite in the sample from protein to which it is bound, to produce a sample comprising unbound vitamin D metabolite and protein; and
- 15 (b) measuring the amount of vitamin D metabolite in the sample of (a).

20 The vitamin D status of a subject is highly informative, and may be used to determine the underlying cause of a number of disease states including rickets, and hyper- or hypo-calcaemia. Thus, in a preferred embodiment of the second aspect of the invention, the method relates to determining the cause of disease in a subject, wherein preferably, the subject is a human and most preferably a child.

25 The present invention is performed *in vitro*, on a sample of serum or plasma removed from a subject. Thus, preferably, the method comprises the additional step of first removing a serum or plasma sample from the subject. The present invention therefore relates to a non-invasive method, the results of which may be used in determining vitamin D status of a subject, and thus the underlying cause of disease. The method does not

however provide a result upon which an immediate medical decision regarding treatment must be made.

5 In a third aspect of the present invention, there is provided a displacement agent as defined in relation to the first aspect.

10 In a fourth aspect of the invention, there is provided the use of an agent of the third aspect. In a method for measuring vitamin D metabolite in a sample of serum or plasma, or for determining the vitamin D status of a subject, preferably the displacement agents are used in methods of the first and second aspects of the invention.

15 In a fifth aspect of the invention, there is provided a kit for use in a method of the first or second aspects, comprising a displacement agent according to the second aspect of the invention. The kit preferably also comprises a key showing the correlation between the results of the assay with the amount of vitamin D metabolite present in the sample. In a further preferred embodiment, a kit may comprise means for carrying out step (b) of the first aspect of the invention. Such means may include one or more of a support, labelled vitamin D metabolites, antibodies, proteins on labels. The kit  
20 preferably will also comprise instructions for use.

The preferred embodiments apply to each aspect, *mutatis mutandis*.

25 The present invention will now be described by way of a non-limiting example, with reference to the following figures, in which:

Figure 1 shows a typical calibration curve for the direct 25-hydroxyvitamin D enzyme immunoassay.

Figure 2 shows a correlation of the direct enzyme immunoassay of the present invention with Immunodiagnostic Systems Ltds' Gamma B 25-Hydroxy Vitamin D radioimmunoassay (IDS catalogue number AA-35F1) with 180 serum or plasma patient samples.

5

Figure 3 shows a correlation of the direct enzyme immunoassay with the Diasorn 25-Hydroxy Vitamin D <sup>125</sup>I RIA (catalogue number 68100E, Diasorin, Stillwater, Minnesota, USA) with 55 serum or plasma patient samples

10

### **Example**

#### **Preparation of anti-vitamin D coated microtitre plates**

Donkey anti-(sheep IgG) serum and sheep anti-vitamin D were both purified by sodium sulphate precipitation to give the corresponding IgG fractions. Microtitre plates (Nunc Maxisorp) were coated with donkey anti(sheep IgG), a purified IgG, 250µl per well of 30mg/l IgG in 10mM phosphate buffered saline (PBS) and incubated overnight at room temperature. The coated plates were washed three times with 10mM phosphate buffered saline, 0.05% Tween 20(PBST). Sheep anti-vitamin IgG, 1ng/ml in PBS, 0.1% Polypep (Sigma), 0.05% sodium azide, 200µl per well was added and plates stored at 2-8C. Plates were washed with PBST before use.

20

#### **Preparation of vitamin D biotin**

A solution of 24,25-dihydroxyvitamin D in methanol (200µl/mg/mL) was treated with an excess of sodium periodate in water for 1 hour at room temperature and purified by HPLC chromatography (Hypersil C18 column 4.6mm x 125mm, 50% methanol/water, elution gradient of 50% to 100% methanol). The eluted fractions containing (25,26,27-nor)-vitamin D-24-aldehyde were treated with excess carboxymethyl oxime (Sigma) methanol for 4 hours, purified by HPLC chromatography (as above) to provide 24-CMO-vitamin D. The N-

25

hydroxysuccinimide ester was prepared by treatment of 24-CMO-vitamin D with excess N- hydroxysuccinimide and dicyclohexylcarbodiimide in dioxan. The N-hydroxysuccinimide ester of 24-CMO-vitamin D and biotin N- hydroxysuccinimide ester (Sigma) were added to bisaminopolyethylene glycol (Sigma) in dioxan solution and allowed to react for 3 hours at room temperature. Excess 24-CMO-vitamin D and biotin were removed by dialysis and the vitamin D-biotin conjugate was stored at -20C.

#### **Vitamin D displacement agent**

The preferred formulation of the vitamin D displacement agent is any conventional buffer, such as phosphate buffered saline solution containing 8-anilino-1-naphthalenesulphonic acid ammonium salt (0.5 to 10g/l, preferably 1.6g/l), 3-( $\alpha$ -acetylbenzyl)-4-hydroxycoumarin (50 to 1000mg/l, preferable 160mg/l) and methanol (10 to 300m/l, preferably 160m/l).

#### **Enzymeimmunoassay**

A typical assay procedure is as follows:

A portion of sample (25 $\mu$ l serum or plasma) is diluted with 1ml of vitamin D displacement agent as described above. A portion of the diluted sample (100 $\mu$ l) is added to the anti-vitamin D antibody coated microtitre plate, followed by addition of a solution (100 $\mu$ l )of the vitamin D-biotin conjugate, and incubated for 90 minutes at room temperature. The plate was washed three times with 10mM phosphate buffered saline containing 0.05% Tween20 (PBST). Avidin peroxidase conjugate (Sigma) diluted 1:2000 in PBST was added and incubated for 30 minutes at room temperature, followed by washing three times with PBST. TMB substrate reagent (Moss Inc.) was added and colour allowed to develop for 30 minutes. After addition of 0.5M HCL to stop the reaction, the absorbance was recorded at 450nm.



A calibration curve is prepared with each batch of samples being analysed, and the 25-hydroxyvitamin D values for each sample can be read directly from the calibration curve using the absorbance value obtained for each sample. This can be performed manually from a calibration curve plotted on graph paper, or more usually by suitable data processing software.

A typical calibration curve for the direct 25-hydroxyvitamin D enzymeimmunoassay is shown in Figure 1.

10

	25-Hydroxyvitamin D ( nmol/L)	Absorbance at 450nm	B/Bo (%)
15	0	2.436	100%
	6.8	2.015	83%
	14	1.678	69%
	27	1.330	55%
	67	0.815	33%
20	179	0.484	20%
	380	0.332	14%

To demonstrate effectiveness of the direct 25-hydroxyvitamin D enzymeimmunoassay with patient samples, the assay was compared to two commercially available radioimmunoassays, both of which use an extraction step as part of sample preparation prior to assay. The two radioimmunoassays are widely used clinically for the quantitative determination of 25-hydroxyvitamin D and other metabolites in human serum or plasma as part of the assessment of vitamin D sufficiency. The direct 25D enzymeimmunoassay shows good agreement and correlation with two established

extraction radioimmunoassays, thus demonstrating the utility of a direct 25-hydroxyvitamin D enzymeimmunoassay for the quantitative determination of 25-hydroxyvitamin D (and other metabolites) in serum or plasma specimens (Figures 2 and 3).

Claims

1. A method for measuring vitamin D metabolite present in a plasma or serum  
5 sample, the method comprising:
- (a) adding to a serum or plasma sample a non-competitive displacement agent to  
effect separation of any vitamin D metabolite in the sample from protein to which it is  
bound, such that any vitamin D metabolite may be detected and/or measured; and  
(b) detecting or measuring the amount of vitamin D metabolite in the sample of  
10 (a).
2. A method for measuring vitamin D metabolite in a serum or plasma sample  
according to claim 1, wherein the vitamin D metabolite is 25-hydroxy Vitamin D or an  
analogue thereof.
- 15 3. A method for measuring vitamin D metabolite in a serum or plasma sample  
according to claim 1 or 2, wherein the non-competitive displacement agent is a  
chemical reagent.
- 20 4. A method for measuring vitamin D metabolite in a serum or plasma sample  
according to claim 3 wherein the displacement agent comprises 8-anilino-1-  
naphthalenesulphonic acid ammonium salt, 3-(acetonylbenzyl)-4-hydroxycoumarin and  
a water miscible solvent.
- 25 5. A method for measuring vitamin D metabolite in a serum or plasma sample  
according to claim 5, wherein the water miscible solvent is methanol.
6. A method for measuring vitamin D metabolite in a serum or plasma sample  
according to any one of claims 1 to 5 wherein the amount of vitamin D metabolite is  
30 measured in a binding assay.

7. A method for measuring vitamin D metabolite in a serum or plasma sample according to claim 6 wherein the binding assay is a competitive binding assay.
- 5 8. A method for measuring vitamin D metabolite in a serum or plasma sample according to claim 7 wherein the competitive binding assay comprises the steps of
- (a) providing a support having immobilised thereon a binding factor capable of binding a vitamin D metabolite;
- (b) contacting the support with a sample comprising the vitamin D metabolite to be  
10 measured;
- (c) contacting the support with a labeled form of the vitamin D metabolite;
- (d) measuring the amount of labeled vitamin D metabolite left bound to the support,  
wherein the amount of labeled vitamin D metabolite bound to the support is  
15 proportional to the amount of vitamin D metabolite in the sample.
9. A method for measuring vitamin D metabolite in a serum or plasma sample according to claim 8, wherein the binding factor immobilised on the support is DBP.
- 20 10. A method for measuring vitamin D metabolite in a serum or plasma sample according to claim 8, wherein the binding factor immobilised on the support is an anti-vitamin D metabolite antibody.
11. A method for measuring vitamin D metabolite in a serum or plasma sample  
25 according to any one of claims 8 to 10, wherein the vitamin D metabolite is labeled with biotin, avidin, a fluorescent molecule, or a chemiluminescent molecule.
12. A method for measuring vitamin D metabolite in a serum or plasma sample according to claim 6, wherein the binding assay is an immunoassay.

13. A method for determining the vitamin D status of a subject, comprising the method according to any one of claims 1 to 12.
14. A method for determining the vitamin D status of a subject according to claim  
5 13 comprising the additional step of first removing a sample from a subject.
15. A non-competitive displacement agent according to any one of claims 3 to 5.
16. Use of a non-competitive displacement agent according to claim 15 in  
10 measuring the vitamin D metabolite in a serum or plasma sample.
17. Use of a non-competitive displacement agent according to claim 15 in a method according to any one of claims 1 to 14.
18. A kit for use in measuring vitamin D metabolite in a sample comprising a  
15 displacement agent according to claim 15.
19. A kit according to claim 18 further comprising a key showing the correlation between the results of the method and the amount of vitamin D metabolite  
20 present in the sample.
20. A kit according to claim 18 or 19 further comprising means for measuring the vitamin D metabolite present in the sample.
21. A kit according to any one of claims 18 to 20 further comprising one or  
25 more of a support, labels, protein, antibody and instructions for use.


Figure 1


Figure 2

Figure 3